

PART 1 - CHEMISTRY

Topic : Electrochemistry

SECTION-A : (Maximum Marks : 12)

- This section contains **THREE** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is(are) correct answer(s).
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme :

Full Marks	: +4	ONLY if (all) the correct option(s) is(are) chosen;
Partial Marks	: +3	If all the four options are correct but ONLY three options are chosen;
Partial Marks	: +2	If three or more options are correct but ONLY two options are chosen, both of which are correct;
Partial Marks	: +1	If two or more options are correct but ONLY one option is chosen and it is a correct option;
Zero Marks	: 0	If none of the options is chosen (i.e. the question is unanswered);
Negative Marks	: -1	In all other cases.
- For example, in a question, if **(A), (B) and (D)** are the ONLY three options corresponding to correct answers, then choosing ONLY (A), (B) and (D) will get +4 marks; choosing ONLY (A) and (B) will get +2 marks; choosing ONLY (A) and (D) will get +2 marks; choosing ONLY (B) and (D) will get +2 marks; choosing ONLY (A) will get +1 mark; choosing ONLY (B) will get +1 mark; choosing ONLY (D) will get +1 mark; choosing no option (i.e. the question is unanswered) will get 0 marks; and choosing any other combination of options will get -1 marks.

1. At 300 K, the molar conductivities of the aqueous solutions of three salts at two different concentrations are given below:

Salt	Concentration (M)	Molar conductivity ($\text{S cm}^2 \text{mol}^{-1}$)
NaNO_3	0.01	111
	0.04	101
NaCl	0.01	117
	0.04	107
AgNO_3	0.01	125
	0.04	116

The conductivity of a saturated aqueous solution of AgCl is $1.40 \times 10^{-6} \text{ S cm}^{-1}$ at 300 K. If the solubility of AgCl in water at 300 K is $X \text{ mol L}^{-1}$, then $\log_{10}(X^{-1})$ is

(Assume that AgCl dissolved in water ionizes completely and that the molar conductivity of saturated AgCl solution is equal to its limiting molar conductivity.)

[JEE Advanced 2026 Paper 2]

- (A) 3 (B) 4 (C) 5 (D) 6

Scan or tap for the
full solution



Space for Rough Work





2. In a conductometric titration, small volume of titrant of higher concentration is added stepwise to a larger volume of titrate of much lower concentration, and the conductance is measured after each addition. The limiting ionic conductivity (Λ_0) values (in $\text{mS m}^2 \text{mol}^{-1}$) for different ions in aqueous solutions are given below:

Ions	Ag^+	K^+	Na^+	H^+	NO_3^-	Cl^-	SO_4^{2-}	OH^-	CH_3COO^-
Λ_0	6.2	7.4	5.0	35.0	7.2	7.6	16.0	19.9	4.1


For different combinations of titrates and titrants given in **List-I**, the graphs of 'conductance' versus 'volume of titrant' are given in **List-II**.

Match each entry in **List-I** with the appropriate entry in **List-II** and choose the correct option.

[JEE Advanced 2024 Paper 1]

List-I		List-II	
P.	Titrate: KCl Titrant: AgNO_3	1.	
Q.	Titrate: AgNO_3 Titrant: KCl	2.	
R.	Titrate: NaOH Titrant: HCl	3.	
S.	Titrate: NaOH Titrant: CH_3COOH	4.	

Space for Rough Work

List-I	List-II
	5. 

- (A) P-4, Q-3, R-2, S-5 (B) P-2, Q-4, R-3, S-1 (C) P-3, Q-4, R-2, S-5 (D) P-4, Q-3, R-2, S-1

Scan or tap for the full solution



3. An aqueous solution of hydrazine (N_2H_4) is electrochemically oxidized by O_2 , thereby releasing chemical energy in the form of electrical energy. One of the products generated from the electrochemical reaction is $\text{N}_2(\text{g})$.

Choose the correct statement(s) about the above process

[JEE Advanced 2024 Paper 2]

- (A) OH^- ions react with N_2H_4 at the anode to form $\text{N}_2(\text{g})$ and water, releasing 4 electrons to the anode.
 (B) At the cathode, N_2H_4 breaks to $\text{N}_2(\text{g})$ and nascent hydrogen released at the electrode reacts with oxygen to form water.
 (C) At the cathode, molecular oxygen gets converted to OH^- .
 (D) Oxides of nitrogen are major by-products of the electrochemical process.

Scan or tap for the full solution



Space for Rough Work

SECTION-B : (Maximum Marks : 8)

- This section contains **TWO** questions.
- The answer to each question is a **NUMERICAL VALUE**.
- For each question, enter the correct numerical value of the answer (in decimal notation, truncated / rounded-off to the **SECOND DECIMAL PLACE**; e.g. 6.25, 7.00, -0.33, -0.30, 30.27, -127.30) in the place designated to enter the answer.
- Answer to each question will be evaluated according to the following marking scheme :
 Full Marks : +4 If **ONLY** the correct answer is entered;
 Zero Marks : 0 In all other cases.

4. In an electrochemical cell, dichromate ions in aqueous acidic medium are reduced to Cr^{3+} . The current (in amperes) that flows through the cell for 48.25 minutes to produce 1 mole of Cr^{3+} is _____.

Use: 1 Faraday = 96500 C mol^{-1}

[JEE Advanced 2025 Paper 1]

Scan or tap for the full solution



5. An electrochemical cell is fueled by the combustion of butane at 1 bar and 298 K. Its cell potential is $\frac{X}{F} \times 10^3$ volts, where F is the Faraday constant. The value of X is _____.

Use: Standard Gibbs energies of formation at 298 K are: $\Delta_f G^\circ_{\text{CO}_2} = -394 \text{ kJ mol}^{-1}$; $\Delta_f G^\circ_{\text{water}} = -237 \text{ kJ mol}^{-1}$; $\Delta_f G^\circ_{\text{butane}} = -18 \text{ kJ mol}^{-1}$

[JEE Advanced 2025 Paper 2]

Scan or tap for the full solution



Space for Rough Work

JEE ADVANCED ELECTROCHEMISTRY — PYQS

Test Pattern : **JEE (Advanced)**

ANSWER KEY

SECTION-A

[illegible]

SECTION-B

Ques.	4	5								
Ans.	100	105.4 to 105.6								