

PART 1 - CHEMISTRY

Topic : States of matter

SECTION-A : (Maximum Marks : 8)

- This section contains **TWO** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is(are) correct answer(s).
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme :

Full Marks	: +4	ONLY if (all) the correct option(s) is(are) chosen;
Partial Marks	: +3	If all the four options are correct but ONLY three options are chosen;
Partial Marks	: +2	If three or more options are correct but ONLY two options are chosen, both of which are correct;
Partial Marks	: +1	If two or more options are correct but ONLY one option is chosen and it is a correct option;
Zero Marks	: 0	If none of the options is chosen (i.e. the question is unanswered);
Negative Marks	: -1	In all other cases.
- For example, in a question, if **(A), (B) and (D)** are the ONLY three options corresponding to correct answers, then choosing ONLY (A), (B) and (D) will get +4 marks; choosing ONLY (A) and (B) will get +2 marks; choosing ONLY (A) and (D) will get +2 marks; choosing ONLY (B) and (D) will get +2 marks; choosing ONLY (A) will get +1 mark; choosing ONLY (B) will get +1 mark; choosing ONLY (D) will get +1 mark; choosing no option (i.e. the question is unanswered) will get 0 marks; and choosing any other combination of options will get -1 marks.

- The correct statement(s) about intermolecular forces is(are) [JEE Advanced 2025 Paper 2]

(A) The potential energy between two point charges approaches zero more rapidly than the potential energy between a point dipole and a point charge as the distance between them approaches infinity.

(B) The average potential energy of two rotating polar molecules that are separated by a distance r has $1/r^3$ dependence.

(C) The dipole-induced dipole average interaction energy is independent of temperature.

(D) Nonpolar molecules attract one another even though neither has a permanent dipole moment.
- A closed vessel contains 10 g of an ideal gas X at 300 K, which exerts 2 atm pressure. At the same temperature, 80 g of another ideal gas Y is added to it and the pressure becomes 6 atm. The ratio of root mean square velocities of X and Y at 300 K is [JEE Advanced 2024 Paper 1]

(A) $2\sqrt{2} : \sqrt{3}$ (B) $2\sqrt{2} : 1$ (C) 1:2 (D) 2:1

Scan or tap for the full solution



Scan or tap for the full solution



Space for Rough Work

SECTION-B : (Maximum Marks : 8)

- This section contains **TWO** questions.
- The answer to each question is a **NUMERICAL VALUE**.
- For each question, enter the correct numerical value of the answer (in decimal notation, truncated / rounded-off to the **SECOND DECIMAL PLACE**; e.g. 6.25, 7.00, -0.33, -0.30, 30.27, -127.30) in the place designated to enter the answer.
- Answer to each question will be evaluated according to the following marking scheme :
 Full Marks : +4 If ONLY the correct answer is entered;
 Zero Marks : 0 In all other cases.

3. Two cylinders, both fitted with frictionless pistons, are filled with mixtures of He and Ar gases. In the first cylinder, the masses of He and Ar are m_1 and m_2 , respectively. In the second cylinder, the masses of He and Ar are m_2 and m_1 , respectively. The molar mass of Ar is 10 times the molar mass of He. The external pressure applied by the piston on the first cylinder needs to be 5 times that on the second cylinder so that the volume of the gas mixtures in both the cylinders are equal at the same temperature. Assuming He and Ar behave like ideal gases, the value of (m_1/m_2) is _____. [JEE Advanced 2026 Paper 1]

Scan or tap for the
full solution



4. Molar volume (V_m) of a van der Waals gas can be calculated by expressing the van der Waals equation as a cubic equation with V_m as the variable. The ratio (in mol dm^{-3}) of the coefficient of V_m^2 to the coefficient of V_m for a gas having van der Waals constants $a = 6.0 \text{ dm}^6 \text{ atm mol}^{-2}$ and $b = 0.060 \text{ dm}^3 \text{ mol}^{-1}$ at 300 K and 300 atm is _____.

Use: Universal gas constant (R) = $0.082 \text{ dm}^3 \text{ atm mol}^{-1} \text{ K}^{-1}$

[JEE Advanced 2025 Paper 1]

Scan or tap for the
full solution



Space for Rough Work

JEE ADVANCED STATES OF MATTER — PYQS

Test Pattern : **JEE (Advanced)**

ANSWER KEY

SECTION-A

[illegible]

SECTION-B

[illegible]