

PART 1 - CHEMISTRY

Topic : Surface chemistry

SECTION-B : (Maximum Marks : 4)

- This section contains **ONE** question.
- The answer to each question is a **NON-NEGATIVE INTEGER**.
- For each question, enter the correct integer corresponding to the answer in the place designated to enter the answer.
- Answer to each question will be evaluated according to the following marking scheme :

Full Marks : +4 If ONLY the correct answer is entered;

Zero Marks : 0 In all other cases.

1. To form a complete monolayer of acetic acid on 1g of charcoal, 100 mL of 0.5 M acetic acid was used. Some of the acetic acid remained unadsorbed. To neutralize the unadsorbed acetic acid, 40 mL of 1 M NaOH solution was required. If each molecule of acetic acid occupies $P \times 10^{-23} \text{ m}^2$ surface area on charcoal, the value of P is _____.

[Use given data: Surface area of charcoal = $1.5 \times 10^2 \text{ m}^2 \text{ g}^{-1}$; Avogadro's number (N_A) = $6.0 \times 10^{23} \text{ mol}^{-1}$]

[JEE Advanced 2024 Paper 2]

Scan or tap for the full solution



Space for Rough Work

**JEE ADVANCED SURFACE CHEMISTRY — ADSORPTION —
PYQS**

Test Pattern : **JEE (Advanced)**

ANSWER KEY

SECTION-B

[illegible]