

NEET NUCLEI — NUCLEAR FISSION — PYQS

Max. Marks: 24
Questions: 6

Section A (MCQ) — 6 questions

Q.1 Water is used as a coolant in a nuclear reactor because of its [RE-NEET (UG) 2024]

- (1) high thermal expansion coefficient
- (2) high specific heat capacity
- (3) low density
- (4) low boiling point

Scan or tap for the full solution



Q.2 A nucleus with mass number 240 breaks into two fragments each of mass number 120, the binding energy per nucleon of unfragmented nuclei is 7.6 MeV while that of fragments is 8.5 MeV. The total gain in the Binding Energy in the process is :

[NEET (UG) 2021]

- (1) 216 MeV
- (2) 0.9 MeV
- (3) 9.4 MeV
- (4) 804 MeV

Scan or tap for the full solution



Q.3 When a uranium isotope $^{235}_{92}\text{U}$ is bombarded with a neutron, it generates $^{89}_{36}\text{Kr}$, three neutrons and :

[NEET (UG) 2020]

- (1) $^{91}_{40}\text{Zr}$
- (2) $^{101}_{36}\text{Kr}$
- (3) $^{103}_{36}\text{Kr}$
- (4) $^{144}_{56}\text{Ba}$

Scan or tap for the full solution



Q.4 In any fission process the ratio $\frac{\text{mass of fission products}}{\text{mass of parent nucleus}}$ is

[AIPMT 2005]

- (1) equal to 1
- (2) greater than 1
- (3) less than 1
- (4) depends on the mass of the parent nucleus.

Scan or tap for the full solution



Q.5 Energy released in nuclear fission is due to

[AIPMT 2001]

- (1) some mass is converted into energy
- (2) total binding energy of fragments is more than the binding energy of parental element
- (3) total binding energy of fragments is less than the binding energy of parental element
- (4) total binding energy of fragments is equal to the binding energy of parental element

Scan or tap for the full solution



Q.6 Nuclear fission is best explained by [AIPMT 2000]

- (1) liquid drop model
- (2) Yukawa π -meson theory
- (3) independent particle model of the nucleus
- (4) proton-proton cycle

Scan or tap for the full solution



ANSWER KEY

Ques.	1	2	3	4	5	6					
Ans.	2	1	4	3	1	1					