

NEET EVOLUTION — MECHANISM OF EVOLUTION, HARDY-WEINBERG PRINCIPLE, GENETIC DRIFT — PYQS

Max. Marks: 108

Questions: 27

Section A (MCQ) — 27 questions

Q.1 A population of diploid organisms is at Hardy-Weinberg equilibrium. If the frequency of allele A is 0.1, the frequency of AA is _____.

[RE-NEET (UG) 2026]

- (1) 0.99 (2) 0.01 (3) 0.02 (4) 0.10

Scan or tap for the full solution



Q.2 Natural selection can lead to _____.

- A. stabilisation B. genetic drift
C. directional change D. disruption

[RE-NEET (UG) 2026]

- (1) A and C only (2) A only
(3) A, C and D only (4) A, B, C and D

Scan or tap for the full solution



Q.3 Which one of the following factors will not affect the Hardy-Weinberg equilibrium?

[NEET (UG) 2024]

- (1) Genetic recombination
(2) Genetic drift
(3) Gene migration
(4) Constant gene pool

Scan or tap for the full solution



Q.4 Match List-I with List-II :

List - I		List - II	
A.	Gene pool	I.	Stable within a generation
B.	Genetic drift	II.	Change in gene frequency by chance
C.	Gene flow	III.	Transfer of genes into or out of population
D.	Gene frequency	IV.	Total number of genes and their alleles

Choose the correct answer from the options given below :

[RE-NEET (UG) 2024]

- (1) A-III, B-II, C-I, D-IV (2) A-IV, B-II, C-III, D-I
(3) A-I, B-II, C-III, D-IV (4) A-II, B-III, C-IV, D-I

Scan or tap for the full solution



Q.5 Natural selection where more individuals acquire specific character value other than the mean character value, leads to

[NEET (UG) 2022]

- (1) Stabilising change (2) Directional change
(3) Disruptive change (4) Random change

Scan or tap for the full solution



Q.6 The factor that leads to Founder effect in a population is :

[NEET (UG) 2021]

- (1) Genetic drift
(2) Natural selection
(3) Genetic recombination
(4) Mutation

Scan or tap for the full solution



Q.7 A gene locus has two alleles A, a. If the frequency of dominant allele A is 0.4, then what will be the frequency of homozygous dominant, heterozygous and homozygous recessive individuals in the population?

[NEET (UG) 2019]

- (1) 0.16 (AA); 0.48 (Aa); 0.36 (aa)
(2) 0.16 (AA); 0.36 (Aa); 0.48 (aa)
(3) 0.36 (AA); 0.48 (Aa); 0.16 (aa)
(4) 0.16 (AA); 0.24 (Aa); 0.36 (aa)

Scan or tap for the full solution



Q.8 In a species, the weight of newborn ranges from 2 to 5 kg. 97% of the newborn with an average weight between 3 to 3.3 kg survive whereas 99% of the infants born with weights from 2 to 2.5 kg or 4.5 to 5 kg die. Which type of selection process is taking place?

[NEET (UG) 2019]

- (1) Disruptive Selection (2) Cyclical Selection
(3) Directional Selection (4) Stabilizing Selection

Scan or tap for the full solution



Q.9 Artificial selection to obtain cows yielding higher milk output represents :

[NEET 2017]

- (1) directional as it pushes the mean of the character in one direction
- (2) stabilizing followed by disruptive as it stabilizes the population to produce higher yielding cows.
- (3) disruptive as it splits the population into two, one yielding higher output and the other lower output.
- (4) stabilizing selection as it stabilizes this character in the population.

Scan or tap for the full solution 

Q.10 Genetic drift operates in [NEET 2016 (Phase 2)]

- (1) slow reproductive population
- (2) large isolated population
- (3) non-reproductive population
- (4) small isolated population

Scan or tap for the full solution 

Q.11 In Hardy-Weinberg equation, the frequency of heterozygous individual is represented by

[NEET 2016 (Phase 2)]

- (1) pq (2) p^2 (3) $2pq$ (4) q^2

Scan or tap for the full solution 

Q.12 The tendency of population to remain in genetic equilibrium may be disturbed by : [NEET 2013]

- (1) lack of random mating
- (2) random mating
- (3) lack of migration
- (4) lack of mutations

Scan or tap for the full solution 

Q.13 Variation in gene frequencies within populations can occur by chance rather than by natural selection. This is referred to as : [NEET 2013]

- (1) Genetic drift (2) Genetic load
(3) Genetic flow (4) Random mating

Scan or tap for the full solution 

Q.14 Random unidirectional change in allele frequencies that occurs by chance in all populations and especially in small populations is known as :-

[NEET 2013 (Karnataka)]

- (1) Genetic drift (2) Migration
(3) Mutation (4) Natural selection

Scan or tap for the full solution 

Q.15 Genetic variation in a population arises due to :

[NEET 2013 (Karnataka)]

- (1) Reproductive isolation and selection
- (2) Mutations only
- (3) Mutations as well as recombination
- (4) Recombination only

Scan or tap for the full solution 

Q.16 If the mean and the median pertaining to a certain character of a population are of the same value, the following is most likely to occur:- [AIPMT 2007]

- (1) A normal distribution
- (2) A T-shaped curve
- (3) A skewed curve
- (4) A bi-modal distribution

Scan or tap for the full solution 

Q.17 Using imprints from a plate with complete medium and carrying bacterial colonies, you can select streptomycin resistant mutants and prove that such mutations do not originate as adaptation. These imprints need to be used - [AIPMT 2005]

- (1) On plates with minimal medium
- (2) Only on plates with streptomycin
- (3) Only on plates without streptomycin
- (4) On plates with and without streptomycin

Scan or tap for the full solution 

Q.18 At a particular locus, frequency of 'A' allele is 0.6 and that of 'a' is 0.4. What would be the frequency of heterozygotes in a random mating population at equilibrium? [AIPMT 2005]

- (1) 0.48 (2) 0.24 (3) 0.36 (4) 0.16

Scan or tap for the full solution 

Q.19 Which of the following is not true for a species ?

[AIPMT 2005]

- (1) Variations occur among members of a species
- (2) Gene flow does not occur between the populations of a species
- (3) Each species is reproductively isolated from every other species
- (4) Members of a species can interbreed

Scan or tap for the full solution 

Q.20 Random genetic drift in a population probably results from [AIPMT 2003]

- (1) Constant low mutation rate
- (2) Interbreeding within this population
- (3) Large population size
- (4) Highly genetically variable individuals

Scan or tap for the
full solution



Q.21 In a random mating population in equilibrium, which of the following brings about a change in gene frequency in a non-directional manner :-

[AIPMT 2003]

- (1) Selection
- (2) Random drift
- (3) Mutations
- (4) Migration

Scan or tap for the
full solution



Q.22 Which of the following is important for speciation ?

[AIPMT 2002]

- (1) Behavioural isolation
- (2) Seasonal isolation
- (3) Tropical isolation
- (4) Reproductive isolation

Scan or tap for the
full solution



Q.23 Genetic drift operates in : -

[AIPMT 2002]

- (1) Slow reproductive population
- (2) Small isolated population
- (3) Large isolated population
- (4) Fast reproductive population

Scan or tap for the
full solution



Q.24 In which condition, the gene ratio remains constant for any species population ?

[AIPMT 2002]

- (1) Gene flow
- (2) Mutation
- (3) Sexual selection
- (4) Random mating

Scan or tap for the
full solution



Q.25 Reason of diversity in living beings : -

[AIPMT 2001]

- (1) Mutation
- (2) Short term evolutionary change
- (3) Long term evolutionary change
- (4) Gradual change

Scan or tap for the
full solution



Q.26 Forthcoming generations are less adaptive than their parental generation due to

[AIPMT 2001]

- (1) Genetic drift
- (2) Natural selection
- (3) Mutation
- (4) Adaptation

Scan or tap for the
full solution



Q.27 In Lederberg's replica plating experiment, what shall be used to obtain a streptomycin-resistant strain?

[AIPMT 2001]

- (1) Only minimal medium
- (2) Complete medium and streptomycin
- (3) Only complete medium
- (4) Minimal medium and streptomycin

Scan or tap for the
full solution



ANSWER KEY

Ques.	1	2	3	4	5	6	7	8	9	10	11
Ans.	2	3	4	2	2	1	1	4	1	4	3
Ques.	12	13	14	15	16	17	18	19	20	21	22
Ans.	1	1	1	3	1	2	1	2	2	2	4
Ques.	23	24	25	26	27						
Ans.	2	4	3	3	4						